x = 13·42   x̄ 12.2.

THELNETHAM OLD (TM.08 78.)	Quadrats (site 1)	Summary (site 1)	Quadrats (site 2)	Summary (site 2)
Alnus glutinosa.	[illegible]	0.15/15	[illegible]	5 /15
Anagallis tenella.	[illegible]	0.30/30	[illegible]	x
Angelica sylvestris.	[illegible]	0.35/35	[illegible]	.
Briza media.	[illegible]	0.10/10	[illegible]	.
Calamagrostis canescens.	[illegible]	.	[illegible]	x
Caltha palustris.	[illegible]	.	[illegible]	0.1[illegible]/[illegible]
Carex diandra.	[illegible]	0.05/ 5	[illegible]	x
C. elata.	[illegible]	.	[illegible]	x
C. lepidocarpa.	[illegible]	x	[illegible]	0.10/10
C. panicea.	[illegible]	0.20/20	[illegible]	0.20/20
C. pulicaris.	[illegible]	.	[illegible]	x
Cirsium dissectum.	[illegible]	0.05/ 5	[illegible]	.
C. palustre.	[illegible]	.	[illegible]	x
Cladium mariscus.	[illegible]	3 /30	[illegible]	16 /100
Dactylorhiza incarnata.	[illegible]	.	[illegible]	x
D. praetermissa.	[illegible]	x	[illegible]	0.05/ 5
Drosera anglica.	[illegible]	x	[illegible]	.
Epipactis palustris.	[illegible]	x	[illegible]	x
Eupatorium cannabinum.	[illegible]	0.15/15	[illegible]	1 /55
Euphrasia sp.	[illegible]	0.05/ 5	[illegible]	.
Filipendula ulmaria.	[illegible]	0.10/10	[illegible]	.
Galium palustre.	[illegible]	.	[illegible]	0.15/15
G. uliginosum.	[illegible]	0.10/10	[illegible]	0.15/15
Gymnadenia conopsea.var.densa	[illegible]	x	[illegible]	.
Hydrocotyle vulgaris.	[illegible]	0.05/ 5	[illegible]	3 /25
Juncus subnodulosus.	[illegible]	4 /100	[illegible]	9 /80
Lotus uliginosus.	[illegible]	x	[illegible]	x
Lythrum salicaria.	[illegible]	.	[illegible]	0.10/10
Mentha aquatica.	[illegible]	x	[illegible]	0.35/35
Molinia caerulea.	[illegible]	11 /95	[illegible]	15 /65
Oenanthe lachenalii.	[illegible]	0.35/35	[illegible]	1 /65
Parnassia palustris.	[illegible]	0.45/45	[illegible]	0.35/35
Pedicularis palustris.	[illegible]	.	[illegible]	x
Phragmites communis.	[illegible]	0.55/55	[illegible]	11 /65
Pinguicula vulgaris.	[illegible]	.	[illegible]	x
Potentilla erecta.	[illegible]	2 /50	[illegible]	.
Sanguisorba officinalis.	[illegible]	0.25/25	[illegible]	0.10/15
Schoenus nigricans.	[illegible]	18 /100	[illegible]	13 /65
Succisa pratensis.	[illegible]	4 /75	[illegible]	0.65/40
Taraxacum paludosum.	[illegible]	.	[illegible]	x
Triglochin palustris.	[illegible]	.	[illegible]	x
Typha angustifolia.	[illegible]	.	[illegible]	1 /15
Valeriana dioica.	[illegible]	3 /95	[illegible]	3 /70
Vicia cracca.	[illegible]	0.05/ 5	[illegible]	0.05/ 5
Thelypteris palustris.	[illegible]	x	[illegible]	.
Acrocladium cuspidatum.	[illegible]	4 /45	[illegible]	13 /65
A. giganteum.	[illegible]	.	[illegible]	1 /15
Brachythecium calcareum.	[illegible]	x	[illegible]	x
Bryum pseudotriquetrum.	[illegible]	0.40/40	[illegible]	0.10/10
Calypogeia muelleriana.	[illegible]	x	[illegible]	.
C. trichomanis.	[illegible]	0.05/ 5	[illegible]	.
Campylium elodes.	[illegible]	x	[illegible]	1 /30
C. protensum.	[illegible]	.	[illegible]	x
C. stellatum.	[illegible]	9 /65	[illegible]	8 /50
Cephalozia bicuspidata.	[illegible]	0.05/ 5	[illegible]	.
Chiloscyphus pallescens.	[illegible]	0.45/20	[illegible]	0.35/10
Cratoneuron commutatum.	[illegible]	0.05/ 5	[illegible]	x
C. falcatum.	[illegible]	.	[illegible]	x
C. filicinum.	[illegible]	.	[illegible]	x
Ctenidium molluscum.	[illegible]	17 /70	[illegible]	.
Drepanocladus intermedius. }	[illegible]	}	[illegible]	}
D. revolvens. }	[illegible]	17 /75	[illegible]	18 /30
D. vernicosus.	[illegible]	.	[illegible]	x
Fissidens adiantoides.	[illegible]	10 /75	[illegible]	6 /65
Lophocolea cuspidata.	[illegible]	.	[illegible]	x
Marchantia polymorpha.	[illegible]	.	[illegible]	x
Mnium pseudopunctatum.	[illegible]	0.10/10	[illegible]	8 /55
M. rugicum.	[illegible]	.	[illegible]	x
M. seligeri.	[illegible]	0.10/10	[illegible]	3 /15
Moerckia flotowiana.	[illegible]	.	[illegible]	x
Pellia fabbroniana.	[illegible]	.	[illegible]	0.10/10
Philonotis calcarea.	[illegible]	0.05/ 5	[illegible]	1 /15
Pseudoscleropodium purum.	[illegible]	.	[illegible]	x
Rhytidiadelphus squarrosus.	[illegible]	0.05/ 5	[illegible]	.
Riccardia multifida.	[illegible]	0.35/35	[illegible]	0.15/15
R. pinguis.	[illegible]	0.30/35	[illegible]	1 /30
R. sinuata.	[illegible]	1 /55	[illegible]	x
Scorpidium scorpioides.	[illegible]	.	[illegible]	x
Sphagnum plumulosum.	[illegible]	x	[illegible]	.

THELMETHAM OLD. (TM.02 78.) IV a III not done EA.I

	6	9	7	14	13	11	14	9	7	8	8	7	13	15		8												I
Alnus glutinosa.	.	.	.	x	I	2	2	.	x	.	x	.	.	.	8 /40	.	.	.	.	.	.	.	.	.	.	x		P
Anagallis tenella.	.	.	.	.	.	.	I	.	.	.	.	.	.	.	0.66/ 7	.	.	.	.	.	.	.	.	.	.	.		.
Angelica sylvestris.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	0.07/ 7	.	.	.	.	.	.	.	.	.	.	x		P
Berula erecta.	.	x	.	.	x	.	x	.	.	.	.	.	x	.	0.26/26	.	x	x	I	.	.	x	.	.	.	I	/40	.
Betula pubescens.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.		.
Calamagrostis canescens.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Caltha palustris.	x	x	.	.	x	.	x	.	.	x	.	x	.	x	0.46/46	.	.	.	.	.	.	.	.	.	.	.		P
Carex acutiformis.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
C. diandra.	.	x	.	x	.	x	.	.	.	.	.	.	.	x	0.26/26	.	.	.	.	.	.	.	.	.	.	.		.
C. elata.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	0.07/ 7	x	.	x	.	.	x	x	x	.	x	0.60/60		P
C. lepidocarpa.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.		.
C. nigra.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	0.07/ 7	.	.	x	.	.	.	.	.	.	.	0.10/10		.
C. panicea.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	0.07/ 7	.	.	.	.	.	.	.	.	.	.	.		.
C. paniculata.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
C. remota.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Cirsium arvense.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x		.
C. palustre.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	I	.	.	.	.	.	I	/10	P
Cladium mariscus.	x	x	.	.	x	.	x	x	x	I	x	I	x	x	2 /73	.	x	.	.	.	x	.	x	x	.	0.40/40		P
Dactylorchis praetermissa.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	0.07/ 7	.	.	.	.	.	.	.	.	.	.	.		.
Epilobium palustre.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	0.13/13	.	.	.	.	.	.	.	.	.	.	.		.
Epipactis palustris.	.	.	.	.	x	.	.	.	.	.	x	.	.	.	0.13/13	.	.	.	.	.	.	.	.	.	.	.		.
Eupatorium cannabinum.	.	.	.	.	.	.	.	.	I	.	x	.	x	.	x	0.03/26	x	I	I	I	x	.	.	I	x	I	4 /60	P
Filipendula ulmaria.	.	.	.	.	.	.	.	.	x	.	.	.	.	I	I /13	x	.	x	.	I	2	.	2	.	I	8 /60		P
Frangula alnus.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Galium palustre.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	0.07/ 7	.	.	.	.	.	.	.	.	.	.	.		.
G. uliginosum.	.	x	.	.	.	.	.	.	.	.	.	.	x	.	0.13/13	.	.	.	.	.	.	.	.	.	.	.		P
Glechoma hederacea.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Hedera helix.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Humulus lupulus.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Hydrocotyle vulgaris.	.	.	.	.	.	x	.	.	.	.	.	.	x	x	0.19/20	.	.	.	.	.	.	.	.	.	.	.		P
Hypericum tetrapterum.	.	.	.	.	.	x	.	.	.	.	.	.	x	x	0.19/20	.	.	.	.	.	.	.	.	.	.	.		.
Iris pseudacorus.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Juncus subnodulosus.	2	I	I	.	I	I	I	2	x	I	I	3	4	4	3 25 /92	.	.	.	.	.	.	.	.	.	.	x		P
Ligustrum vulgare.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Lonicera periclymenum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Lotus uliginosus.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	0.07/ 7	.	.	.	.	.	.	.	.	.	.	.		.
Lychnis flos-cuculi.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	0.07/ 7	.	.	.	.	.	.	.	.	.	.	.		.
Lycopus europaeus.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Lythrum salicaria.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x		P
Mentha aquatica.	x	.	.	.	.	x	.	x	.	x	x	x	.	.	0.39/40	x	x	.	x	.	.	x	x	.	x	0.60/60		P
Molinia caerulea.	.	.	x	.	.	x	x	.	x	.	.	.	.	.	0.26/26	.	.	.	.	.	.	.	.	.	.	.		.
Oenanthe fistulosa.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	0.07/ 7	.	.	.	.	.	.	.	.	.	.	.		.
O. lachenalii.	.	.	.	x	.	x	x	x	.	.	.	.	x	x	0.39/40	.	.	.	.	.	.	.	.	.	.	x		.
Parnassia palustris.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.		.
Pedicularis palustris.	.	.	.	x	.	.	.	.	.	.	.	.	x	.	0.13/13	.	.	.	.	.	.	.	.	.	.	.		.
Phragmites communis.	x	.	x	I	.	.	x	x	4	3	4	2	I	x	x 16 /79	5	5	4	4	5	4	5	3	3	2	75 /100		P
Pinguicula vulgaris.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.		.
Ranunculus flammula.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	0.07/ 7	.	.	.	.	.	.	.	.	.	.	.		P
Salix cinerea.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Sanguisorba officinalis.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Solanum dulcamara.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	x	.	.	x	.	x	.	0.40/40		P
Succisa pratensis.	.	.	x	x	x	.	.	.	.	.	.	.	.	x	0.26/26	.	.	.	.	.	.	.	.	.	.	.		.
Typha angustifolia.	.	.	.	.	.	.	.	3	.	x	I	2	.	I	7 /40	.	.	.	.	.	.	.	.	.	.	.		.
T. latifolia.	.	x	.	.	x	.	.	.	.	.	.	.	.	.	0.13/13	.	.	.	.	.	.	.	.	.	.	.		P
Urtica dioica.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Valeriana dioica.	.	.	x	x	x	.	x	x	.	.	.	.	x	.	.	0.39/40	.	.	.	x	.	.	.	.	.	0.10/10		.
V. officinalis.	.	.	.	.	.	.	.	x	.	.	.	.	x	.	.	0.13/13	.	.	.	.	.	.	.	.	.	.		P
Vicia cracca.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	I	.	.	.	.	.	I	/10	.
Equisetum arvense.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
E. palustre.	.	.	.	.	x	.	x	.	x	.	.	x	x	.	x	0.39/40	.	.	.	.	.	.	.	.	.	.		.
Acrocladium cuspidatum.	x	.	.	.	.	.	2	.	.	.	.	.	2	.	4 /20	.	.	.	.	.	.	x	.	x	.	0.20/20		.
A. giganteum.	.	.	.	.	.	.	3	.	.	.	.	.	.	.	3 / 7	.	.	.	.	.	.	.	.	.	.	.		.
Brachythecium rivulare.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	x		.
B. rutabulum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	x	.	.	.	0.10/10		.
Bryum pseudotriquetrum.	.	.	.	x	.	.	x	.	.	.	.	.	.	.	0.13/13	.	.	.	.	.	.	.	.	.	.	.		.
Campylium elodes.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.		.
Cephalozia bicuspidata.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Cratoneuron filicinum.	.	.	.	2	.	.	.	.	.	x	.	.	.	.	2 /13	.	.	.	.	.	.	.	.	.	.	.		.
Ctenidium molluscum.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	0.07/ 7	.	.	.	.	.	.	.	.	.	.	.		.
Drepanocladus intermedius.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	0.07/ 7	.	.	.	.	.	.	.	.	.	.	x		.
D. revolvens.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	0.07/ 7	.	.	.	.	.	.	.	.	.	.	.		.
Eurhynchium praelongum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x		P
Lophocolea heterophylla.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		P
Marchantia polymorpha.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x		.
Mnium pseudopunctatum.	.	I	x	.	.	.	.	.	.	.	.	.	.	x	0.79/20	.	.	.	.	.	.	.	.	.	.	.		.
M. punctatum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		.
M. seligeri.	.	I	.	.	.	.	x	.	.	.	.	.	I	.	I /20	.	.	.	.	.	.	.	.	.	.	x		P
M. undulatum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		.
Pellia fabbroniana.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	0.07/ 7	.	.	.	.	.	.	.	.	.	.	.		P
Rhizomnium punctatum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		.
Riccardia multifida.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	x		.
R. pinguis.	.	.	.	I	.	.	.	.	I	.	x	.	.	.	1 /20	.	.	.	.	.	.	.	.	.	.	.		.
Chara sp.																												

[illegible] (T.L. [illegible])

Species	I	II
Agrostis canina var. stolonifera.	x	x
Alnus glut. moss.	.	..5/ 8
Anagallis tenella.	0.08/ 4	+
Angelica sylvestris.	0.04/ 4	+
Anthoxanthum odoratum.	.	x
Apium nodiflorum.	x	x
Berula erecta.	.	..1./1.
Briza media.	0.08/ 4	+
Cardamine pratensis.	.	..03/ 8
Carex elata.	x	+
C. flacca.	x	x
C. lepidocarpa.	0.08/16	..5./ 8
C. panicea.	8 /64	4 /45
C. pulicaris.	.	..1./1.
Cirsium dissectum.	.	..8/ 5
C. palustre.	0.04/ 4	x
Cladium mariscus.	[illegible]/48	3 /65
Dactylorchis praetermissa.	x	.. 3/ 8
D. traunsteineri.	x	.
Epilobium palustre.	[illegible]	.
[illegible] palustris.	x	..2/ 3
Eupatorium cannabinum.	..40 [illegible]	..20/20
Galium palustre.	.	x
G. uliginosum.	[illegible]	0.24/48
Gymnadenia conopsea var. densiflora.	.	..8/ 2
Hydrocotyle vulgaris.	.	..0/ 5
Juncus subnodulosus.	[illegible]/32	30 /1..
Lemna minor.	x	x
L. trisulca.	x	+
Lotus uliginosus.	..4/ 4	x /20
Lychnis flos-cuculi.	x	..5/ 8
Lythrum salicaria.	x	+
Mentha aquatica.	[illegible]	[illegible]
Molinia caerulea.	[illegible]	[illegible] /45
Parnassia palustris.	x	x
Pedicularis palustris.	..4 4	.
Phragmites communis.	[illegible]	. /75
Pinguicula vulgaris.	[illegible]	[illegible] 8
Potentilla erecta.	[illegible]	[illegible]
Rubus fruticosus.	.	x
Schoenus nigricans.	0 /32	.4 /35
Succisa pratensis.	..4 [illegible]	[illegible]/35
Valeriana dioica.	[illegible] 4	. /55
Equisetum arvense.	x	+
E. palustre.	x	..5/ 6
Acrocladium cuspidatum.	6 /24	.6 /56
A. giganteum.	x	+
[illegible]	..4/[illegible]	0.2./2.
Campylium [illegible]	.	x
C. protensum.	.	x
C. stellatum.	46	0.05/..
Cratoneuron commutatum [illegible]	x	+
C. falcatum.	x	x
C. filicinum.	[illegible]	x
Ctenidium molluscum.	3 [illegible]	0.05/ 5
Drepanocladus [illegible]	x	.
D. revolvens var. intermedius.	[illegible]	2 /18
Fissidens adianthoides.	x	0.50/10
[illegible]	..4 4	0.20/20
Mnium pseudopunctatum.	.	0.1./10
M. seligeri.	.	0.55/10
[illegible]	x	+
[illegible]	x	3 /20
[illegible]	0.04 4	.
[illegible]	x	+
Riccardia multifida.	x	+
[illegible]	[illegible]	0.16/16
[illegible]	0.03/12	x
Scorpidium scorpioides.	x	+
[illegible]	x	+
Chara sp.	7 /40	0.08/ 8

Agrostis canina var.stolon.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	x	.	.	0.10/10	.	.	.	.	.	.	.	.	.	.	.	.	.
Anthoxanthum odoratum.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	0.05/ 5	.	.	.	.	.	.	.	x	.	.	0.10/10		
Apium nodiflorum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Berula erecta.	3	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	I	.	2	x	5 /25	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Blysmus compressus.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	0.10/10			
Caltha palustris.	.	.	.	.	.	.	.	.	.	.	.	.	.	I	.	.	.	.	.	.	.	0.50/ 5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Cardamine pratensis.	.	.	.	.	.	.	x	.	.	.	x	.	.	.	.	.	.	.	.	.	.	0.10/10	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Carex appropinquata.	.	.	.	.	.	.	.	I	.	.	.	.	.	.	.	.	.	.	.	.	.	0.50/ 5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
C. oiandra.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	0.05/ 5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
C. lepidocarpa.	.	.	x	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	0.10/10	.	.	.	.	.	.	.	.	.	.	.	.	.	.
C. panicea.	.	x	.	x	x	I	.	.	.	x	.	.	.	.	.	x	I	.	x	I	.	2 /45	.	.	.	.	.	.	.	.	.	x	x	0.20/20		
C. paniculata.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.
C. pulicaris.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	
Cirsium dissectum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.
C. palustre.	.	.	.	I	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	0.55/1.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Cladium mariscus.	.	I	x	5	.	x	I	5	4	.	5	I	3	2	3	2	I	I	x	I		28 /85	I	.	x	.	x	x	x	x	x	.	0.0 0/70			
Drosera rotundifolia.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	x	.	x	0.30/30			
Epipactis palustris.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	0.10/10			
Eupatorium cannabinum.	.	x	.	.	.	.	.	.	x	.	.	.	.	.	.	.	x	.	.	.	.	0.15/15	.	.	x	.	.	.	.	.	.	.	0.15/15			
Galium palustre.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.
G. uliginosum.	.	.	.	x	x	.	.	.	.	x	.	.	.	.	.	.	.	.	x	.	.	0.20/20	.	.	.	x	.	.	.	x	x	.	0.30/30			
Gymnadenia conopsea var.cons.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	0.05/ 5	.	.	x	.	.	x	.	.	.	x	0.30/30			
Holcus lanatus.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	x	.	.	x	.	.	.	.	.	0.20/20			
Juncus subnodulosus.	x	I	3	.	2	3	x	x	.	3	x	3	2	.	2	.	3	3	2	4		16 /80	I	I	2	2	I	x	I	x	2	I	14 /100			
Lotus uliginosus.	.	.	.	x	x	I	.	.	.	.	.	.	x	.	x	.	.	x	.	.	.	0.75/30	x	.	.	x	.	I	x	I	.	x	2 /60			
Lychnis flos-cuculi.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	0.05/ 5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Mentha aquatica.	x	x	.	x	x	x	.	.	x	x	.	x	.	x	.	x	.	.	.	x	.	0.55/55	.	.	.	.	x	x	.	.	x	.	0.30/30			
Molinia caerulea.	.	.	.	.	x	I	.	.	I	.	x	.	.	x	.	.	.	x	.	.	.	1 /30	x	I	.	I	I	I	x	I	.	x	5 /50			
Parnassia palustris.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Phragmites communis.	x	.	I	.	.	x	x	x	x	I	x	x	.	.	I	.	.	I	x	.	.	2 /60	.	x	x	.	.	x	x	.	x	.	0.50/50			
Potentilla erecta.	.	.	.	.	.	.	.	.	.	x	.	x	.	.	x	.	.	.	.	.	.	0.15/15	x	x	x	x	x	I	I	x	x	x	3 /100			
Schoenus nigricans.	x	3	x	I	.	.	I	.	.	.	.	.	.	I	x	.	.	.	.	.	.	4 /35	.	.	.	x	.	.	.	.	.	x	0.20/20			
Succisa pratensis.	.	.	.	x	.	.	.	.	.	x	.	x	.	.	x	.	.	x	.	.	.	0.25/25	x	x	x	x	x	I	x	I	.	.	3 /50			
Valeriana dioica.	.	x	x	x	x	x	.	.	x	.	.	I	.	.	x	.	.	x	.	.	.	0.90/45	.	.	.	I	x	x	.	.	x	x	1 /50			
Vicia cracca.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	0.05/ 5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Equisetum palustre.	x	.	x	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	0.20/20	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Acrocladium cuspidatum.	.	.	2	.	2	.	.	.	.	.	.	I	.	.	I	I	.	.	.	.	.	5 /25	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Aulacomnium palustre.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.
Campylium protensum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.
C. stellatum.	.	.	I	.	I	.	I	.	.	.	.	.	.	.	x	.	.	.	I	.	.	2 /25	.	.	.	.	.	.	.	.	x	.	0.10/10			
Cratoneuron filicinum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	4	.	.	4 /10	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Drepanoclados intermedius.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	0.05/ 5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Eurhynchium praelongum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Fissidens adiantoides.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	0.05/ 5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Lophocolea cuspidata.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Mnium pseudopunctatum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	0.05/ 5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
M. seligeri.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	0.05/ 5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
M. undulatum.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	0.05/ 5	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Philonotis calcarea.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Pohlia nutans.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	x	.
Pseudoscleropodium purum.	.	.	.	2	.	4	.	.	.	I	.	.	.	.	.	.	.	I	.	.	.	6 /20	.	.	.	x	.	.	.	.	.	I	1 /20			
Sphagnum plumulosum.	.	.	.	.	.	.	.	.	.	.	.	3	.	.	4	.	.	.	.	.	.	6 /10	5	4	5	5	5	5	4	5	4	3	80 /100			
Chara sp.	.	2	2	.	.	.	.	.	.	I	.	.	x	.	.	x	.	.	.	.	.	4 /45	.	.	.	.	.	.	.	.	.	.	.	.	.	.

REDGRAVE (TM.04 78.) 14.3.

Species	M		[illegible]	
Anagallis tenella.	. . . .	x	. . . .	x
Carex dioica.	. . . .	.	. . . .	x
C. elata.	. . . .	.	. . . .	x
C. lepidocarpa.	. . . .	x	. . . .	.
C. nigra.	. . . .	.	. . . .	x
C. panicea.	. . . .	x	. . . .	.
Cirsium dissectum.	. . . .	.	. . . . 2 . . 1 . 1 . . . . x . .	2 /20
C. palustre.	. . . .	.	x x	0.10/10
Cladium mariscus.	x x 1 . . x 1 x x x . x 1 . x . x x x .	3 /70	1 x . 1 1 x . . . . 1 . . 1 . . .	3 /35
Dactylorchis incarnata.	. . . .	.	. . . .	x
D. praetermissa.	. . . .	x	. . . .	x
D. traunsteineri.	. . . .	.	. . . .	x
Drosera anglica.	. . . .	x	. . . .	.
Epipactis palustris.	. . . .	x	. . . .	x
Eupatorium cannabinum.	. . . .	.	x x x 1 1 2 2 . x 2 1 1 x x 1 x . . x 1	6 /65
Galium palustre.	. . . .	.	. . . .	x
G. uliginosum.	. . . .	.	. . . .	x
Gymnadenia conopsea.var.dens.	. . . .	x	. . . .	.
Hydrocotyle vulgaris.	. . . .	.	. . . . . x	0.05/ 5
Juncus subnodulosus.	x x . . . x x . 1 1 1 1 x x 1 1 . 1 x .	4 /70	1 1 x 1 1 x . 2 2 1 x x x 1 . . x x . 1	9 /60
Mentha aquatica.	. . . .	x	. . . . 1 x . . x	0.60/15
Molinia caerulea.	1 x . 1 . 2 . 2 . 1 . . 2 2 . 1 x 3 x 1	11 /65	. 2 2 1 1 1 x . . . 2 2 3 x . . x x x 1	11 /70
Parnassia palustris.	. . . .	x	. . . .	.
Phragmites communis.	. . . x . x . . . . . . x 1 x . . x . x	0.80/35	4 2 2 3 2 3 3 2 2 3 2 2 2 3 3 3 2 3 4 2	43 /100
Pinguicula vulgaris.	. . . .	x	. . . .	.
Potentilla erecta.	. . . .	.	. x . . . x . . . . x . . . x	0.20/20
Schoenus nigricans.	4 3 4 5 4 3 4 3 4 3 4 4 3 3 2 2 3 2 3 4	[illegible]7 /100	x . . . . x . . x . 1 . . . x	0.70/20
Succisa pratensis.	. . . . . . x x . . x . x	0.20/20	. x . . . . x	0.10/10
Valeriana dioica.	. . . .	x	1 x 1 x 1 1 1 . 1 . . 1 1 x x 1 1 1 1 1	8 /100
Equisetum palustre.	. . . .	x	. . . .	.
Acrocladium cuspidatum.	. . . .	.	. 2 2	3 /10
Brachythecium rivulare.	. . . .	x	. . . .	.
B. rutabulum.	. . . .	.	. . . .	x
B. salebrosum.	. . . .	.	. . . .	x
Bryum pseudotriquetrum.	. . . . . x . x . . x . . . . . x . . .	0.20/20	. . . . x . . . . x . . .	0.10/10
Calypogeia trichomanis.	. . . .	.	. . . x . . . . . x . . 2	2 /15
Campylium elodes.	. 1 . 1 . . . . 1 x 1 1	3 /30	. . . . . . x x . 1 1 1 . . . x x . . .	2 /35
C. protensum.	. . . .	.	. . . .	x
C. stellatum.	1 1 3 4 2 2 2 2 2 . 3 2 . 2 2 1 . x 2 3	26 /[illegible]5	3 x x 1 x x 1 2 2 1 . . x x . 1 1 x . 1	9 /80
Chiloscyphus pallescens.	. . . .	.	. . . .	x
Cratoneuron falcatum.	. . . . . x	0.05/ 5	. . . . x	0.05/ 5
Ctenidium molluscum.	. . . x . 2 . x . 1 . 1 2 . . . . 2 . x	6 /40	. 1 1 2 1 x x 1 . . . 2 . . . . . . 3 x	8 /50
Drepanocladus intermedius.	1	1 /10	. . . . 1	0.50/ 5
Fissidens adiantoides.	. . x . . x . . 1 . 1 2 1 . . . 2 . . .	5 /35	. x 2 2 3 1 x 1 2 . 2 1 2 1 1 1 2 x x 1	16 /95
Lophocolea cuspidata.	. . . .	.	. x x x x 1 . . . . 3 x	3 /35
Mnium pseudopunctatum.	. . . .	.	. . . . x . 1 . . 2	2 /15
Moerckia flotoviana.	. . . . x . . . . . x 1	0.60/15	. . . . x	0.05/ 5
Pellia fabroniana.	. . . .	.	. . . . 2 2 . . . . 1 . .	2 /15
Philonotis calcarea.	. . . .	.	. . . .	x
Pleurozium schreberi.	. . . .	.	. . . .	x
Preissia quadrata.	. . . . 1 . x . . . . . 1 . . .	2 /15	. . . .	.
Pseudoscleropodium purum.	. . . .	.	. . . .	x
Riccardia multifida.	x . . . . x . x x . x . . . x . x . . x	0.40/40	. . . .	x
R. pinguis.	. . . x . x x . . . . x x 1 x 2	2 /40	. . . .	x
R. sinuata.	. x . . . . x . . . x x . . . . . . x .	0.25/25	. . . . . . x . 1	1 /10
Scorpidium scorpioides.	x . . . 2 x . x	2 /20	. . . .	.

Species	Group 1	Group 2
Anagallis tenella.	.	0.58/10
Angelica sylvestris.	x	.
Betula pubescens.	.	5 /20
Briza media.	.	0.26/25
Calluna vulgaris.	.	23 /90
Caltha palustris.	x	.
Carex elata.	x	.
C. lepidocarpa.	x	.
C. panicea.	x	0.20/20
C. pulicaris.	.	0.26/25
Cirsium palustre.	[illegible]	0.05/ 5
Cladium mariscus.	24 /100	0.25/25
Dactylorchis praetermissa.	x	.
Drosera anglica.	x	.
D. rotundifolia.	x	0.35/35
Epipactis palustris.	x	0.10/10
Erica tetralix.	.	4 /25
Eupatorium cannabinum.	[illegible] /5	0.05/ 5
Festuca ovina.	.	0.15/15
Galium uliginosum.	1 /50	.
Gymnadenia conopsea var. dens	x	.
Hydrocotyle vulgaris.	0.35/10	.
Juncus subnodulosus.	7 /100	1 /60
Leontodon hispidus.	.	1 /55
L. leysseri.	.	x
Lotus uliginosus.	x	.
Luzula multiflora.	.	0.05/ 5
Lythrum salicaria.	x	.
Mentha aquatica.	0.1[illegible]/10	.
Molinia caerulea.	6 /50	16 /95
Parnassia palustris.	.	0.05/ 5
Pedicularis palustris.	x	0.05/ 5
Phragmites communis.	. /35	0.80/80
Pinguicula vulgaris.	x	.
Potentilla erecta.	x	4 /60
Schoenus nigricans.	0.65/20	0.38/38
Succisa pratensis.	x	7 /90
Thalictrum flavum.	x	.
Ulex europaeus.	.	0.05/ 5
Valeriana dioica.	x	0.40/40
V. officinalis.	0.65/20	.
Equisetum arvense.	x	x
E. palustre.	0.30/3[illegible]	.
Acrocladium cuspidatum.	x	2 /15
Aulacomnium palustre.	x	0.6[illegible]/15
Brachythecium rutabulum.	x	.
Bryum pseudotriquetrum.	0.15/15	.
Calypogeia muelleriana.	.	0.10/10
C. sphagnicola.	.	x
C. trichomanis.	.	0.70/25
Campylium elodes.	. /20	.
C. stellatum.	. /75	0.30/20
Cephalozia bicuspidata.	0.05/ 5	0.05/ 5
C. connivens.	.	0.05/ 5
Chiloscyphus pallescens.	x	.
Cratoneuron falcatum.	. /60	.
C. filicinum.	x	.
Ctenidium molluscum.	4 /65	0.05/ 5
Drepanocladus revolvens. }	0.35/10	.
D. intermedius.	.	0.05/ 5
Fissidens adiantoides.	. /45	0.10/10
Funaria hygrometrica.	.	x
Hypnum cupressiforme.	.	0.05/10
Lophocolea heterophylla.	.	0.05/ 5
Mnium pseudopunctatum.	[illegible]	.
M. punctatum.	x	.
M. seligeri.	x	.
Philonotis calcarea.	x	.
Pseudoscleropodium purum.	x	.
Riccardia latifrons.	.	x
R. multifida.	[illegible]	x
R. pinguis.	[illegible]	.
R. sinuata.	[illegible]	.
Scorpidium scorpioides.	[illegible]	.
Sphagnum plumulosum.	.	0.10/10
S. russowii.	.	0.55/40
Weissia controversa.	.	x

BELGRAVE (Th. 04 7?.)				LA.3.
			R	
Betula pubescens.	. . . .	.	x . x . x . . . 3 I . . x 4 . . 2 5 I .	14 /55
B. verrucosa.	. . . .	.	. . . .	x
Briza media.	. x x . . x . . x . . . . x . . x . . .	[illegible] 35	. . . .	.
Calluna vulgaris.	I x . 2 I I 2 . x . . 2 . . 2 . x 4 . .	[illegible]	. . . x . x . x . . . . x . . x .	0.25/25
Oarex flacca.	. . . . x	[illegible]	. . . .	.
C. panicea.	. . . .	.	. . . .	x
C. pulicaris.	. . . .	x	. . . .	.
Cirsium dissectum.	. x . . . . x	[illegible]	. . . .	x
Dactylorchis praetermissa.	. . . .	x	. . . .	.
Drosera rotundifolia.	x x x . x . x . x . x I . x I x x . x .	[illegible] /7[illegible]	. . x	0.55/10
Epipactis palustris.	. . . . I . . x . x . . x . . x . . . I	I /3[illegible]	. . . .	.
Erica tetralix.	2 . I 2 2 . . 3 . . . . 2 . 2 2 .	12 /40	. x . . x I	0.60/15
Festuca ovina.	1 x x 1 x I x . 2 I . . x x . . 2 . . x	6 /7[illegible]	. . . . . . x . . x . . . x x . . x . .	0.3[illegible]/30
Galium harcynicum.	. . x . . . . x	0.1 /[illegible]	. . . .	.
Genista anglica.	x . . x . . . . x . . . . x . . .	[illegible]	. . . .	.
Gymnadenia conopsea var.dens.	. . I . . . . x . . . . . x .	[illegible]/15	. . . .	.
Juncus subnodulosus.	I I x I I x . x 2 . . . 2 I . x . I . x	6 /30	x . . x . x x x . . . . . . x . . . x x	0.95/50
Leontodon hispidus.	. x x . x . . . x . . . . x . . . . x .	0.35/35	. . . .	.
Lotus uliginosus.	x . . I . . . . I	2 /30	. . . .	.
Luzula multiflora.	x . x . . . . . x x . . . x . . x . . .	0.[illegible]/30	. . . .	.
Molinia caerulea.	. . 2 x I 2 I I x I x I . I . . x I . .	9 /85	2 4 . . 2 I 2 I 2 4 I 3 I x . 2 x x 2 I	23 /95
Parnassia palustris.	. . . . . x x	0.1 /1[illegible]	. . x . . . x . . . . . x x	0.20/[illegible]
Pedicularis palustris.	x . . . . x x . x . . . . . . x .	0.25/25	. . . .	.
Phragmites communis.	. . . x . . . . x x	0.15/15	. . 2 x x x . x x x I 2 I . x x . . I 2	7 /75
Pinguicula vulgaris.	. . x I	[illegible] /10	. . . .	.
Polygala serpyllifolia.	. . . . x . . . . x .	[illegible]	. . . . x	0.05/ 5
Potentilla erecta.	x I x I x x x . x I x x x . . . x x . .	3 /75	x x x . . x x I . x . . . x . x . . x .	0.95/50
Schoenus nigricans.	. . . . . x . . x . . . x . . . x . . x	0.70/25	. . . .	.
Succisa pratensis.	. x x x I . . x . x 2 2 . I x x I . x .	7 /30	. . . .	.
Ulex europaeus.	. x . . . . x	0.10/10	. . . .	.
Valeriana dioica.	. x . . . x . . x . . . . x . . I	2 /30	. . . .	.
Atrichum undulatum.	. . . .	.	. . . .	x
Aulacomnium palustre.	x . . . . . . x . . . . x . . I .	0.65/2[illegible]	. . . . I . . x . I I . . . . x .	2 /25
Bryum pseudotriquetrum.	. . . . x . . . . x	0.10/10	. . . .	.
Calypogeia sphagnicola.	. . x . . x x x x x . . . x . x x . . .	0.45/45	x . . . x . . . . x	0.15/15
C. trichomanis.	. . . . x . . . . . . x x	0.15/15	. . . x . . . . x	0.10/10
Campylium elodes.	. . . .	.	. x x	0.10/10
C. stellatum.	. . . . . x x . . . . . x I	0.65/20	. x x	0.10/10
Campylopus pyriformis.	. . . .	x	. . . .	.
Cephalozia bicuspidata.	. . . . . x	0.05/ 5	. . . .	.
C. connivens.	. x . x . . x I . . . . . x . x	0.75/30	I . . . . x . .	0.55/10
Ctenidium molluscum.	. . x . . x . . . . . . x I	0.65/20	. . . .	.
Dicranella heteromalla.	. . . .	x	. . . .	x
Dicranum scoparium.	. . . .	x	. . . .	.
Encalypta streptocarpa.	. . . .	.	. . . .	x
Hypnum cupressiforme.	. x x . . x . x . . . . . x . . . I . .	0.75/30	. . . .	.
Lepidozia setacea.	. . . .	.	. . . . . x	0.05/ 5
Lophocolea cuspidata.	. . . .	.	. x . . . . x	0.10/10
Lophozia ventricosa.	. . . .	x	. . 3	9 / 5
Mnium pseudopunctatum.	. . x x	0.10/10	. . . .	.
Pellia epiphylla.	. . . .	x	. . . .	.
Pleurozium schreberi.	. . . .	x	. . . .	.
Pohlia nutans.	. . . .	x	. . . .	x
Polytrichum commune.	. . . .	x	. . x . . . . I .	0.55/10
P. juniperinum.	. . . .	x	. . . .	x
Riccardia latifrons.	. . x . . . . x	0.10/10	. . . .	.
Sphagnum fimbriatum.	. . . .	.	. . . 3 2 . 2 . . . 4 . 5 3 2 . . I . .	18 /40
S. palustre.	. . . .	.	. . 5 . I 4 . 3 3 . . . . I 2 4 5 . . .	24 /25
S. plumulosum.	x x / x x . x 2 3 . 3 4 . x 4 2 3 x x .	23 /75	3 I . . . . I . . . . x I	4 /25